

Math 1B – Calculus – A few divergent integrals worked through – Spring '10

1. Explain why the Integral $\int_0^3 \frac{1}{(x-1)^2(x-2)^2} dx$ is improper and then evaluate it.

SOLN: There are vertical asymptotes at $x = 1$ and $x = 2$ which are interior to the interval of integration.

$$\int_0^3 \frac{1}{(x-1)^2(x-2)^2} dx = \int_0^{3/2} \frac{1}{(x-1)^2(x-2)^2} dx + \int_{3/2}^3 \frac{1}{(x-1)^2(x-2)^2} dx$$


To simplify matters, use symmetry and substitute $u = x - 3/2$ whence the integral becomes

$$\int_{-3/2}^{3/2} \frac{1}{(u+1/2)^2(x-1/2)^2} dx = \int_{-3/2}^0 \frac{1}{(u+1/2)^2(x-1/2)^2} dx + \int_0^{3/2} \frac{1}{(u+1/2)^2(x-1/2)^2} dx$$

and then substitute $v = -u$ into the first one so that it is identical to the second and

$$\begin{aligned} \int_{-3/2}^{3/2} \frac{1}{(u+1/2)^2(x-1/2)^2} dx &= 2 \int_0^{3/2} \frac{1}{(u+1/2)^2(x-1/2)^2} dx \\ &= 2 \int_0^{3/2} \frac{2}{u+1/2} + \frac{1}{(u+1/2)^2} - \frac{2}{x-1/2} + \frac{1}{(x-1/2)^2} dx \\ &= 4 \ln(4) + 3 - 4 \lim_{b \rightarrow 1/2^-} \int_0^b \frac{dx}{x-1/2} + 4 \lim_{b \rightarrow 1/2^+} \int_{3/2}^b \frac{dx}{x-1/2} \\ &\quad + 2 \lim_{b \rightarrow 1/2^-} \int_0^b \frac{dx}{(x-1/2)^2} + 2 \lim_{b \rightarrow 1/2^+} \int_b^{3/2} \frac{dx}{(x-1/2)^2} \\ &= 4 \ln(4) + 3 - 4 \lim_{b \rightarrow 1/2^-} \ln \left| b - \frac{1}{2} \right| + 4 \ln \left| \frac{1}{2} \right| + 4 \lim_{b \rightarrow 1/2^+} \ln \left| b - \frac{1}{2} \right| \\ &\quad - \lim_{b \rightarrow 1/2^-} \frac{2}{b-1/2} - 4 + \lim_{b \rightarrow 1/2^+} \frac{2}{b-1/2} - 2 \\ &= 4 \ln(2) + 3 - 4 \lim_{\varepsilon \rightarrow 0} \ln |\varepsilon| + 4 \lim_{\varepsilon \rightarrow 0} \ln |\varepsilon| \\ &\quad \infty - 4 + \infty - 2 \\ &= 4 \lim_{\varepsilon \rightarrow 0} \ln |1| + \infty = \infty \end{aligned}$$

So, while your first intuition about this integral may be that it's going to go like $1/x^p$ around a vertical asymptote where $p = 2$ would converge...it actually diverges. This is more obvious in the partial fractions expansion.



2. Explain why the Integral $\int_0^3 \frac{1}{\sqrt{|x-1|}\sqrt{|x-2|}} dx$ is improper and then evaluate it.

SOLN: There are vertical asymptotes at $x = 1$ and $x = 2$ which are interior to the interval of integration. To simplify matters, use symmetry and substitute $u = x - 3/2$ whence the integral becomes

$$\int_0^3 \frac{1}{\sqrt{x-1}\sqrt{x-2}} dx = \int_{-3/2}^{3/2} \frac{1}{\sqrt{u-1/2}\sqrt{u+1/2}} du = \int_{-3/2}^{3/2} \frac{1}{\sqrt{u^2-1/4}} du = \int_{-3/2}^{3/2} \frac{2}{\sqrt{4u^2-1}} du$$

Now use y-axis symmetry and then split the absolute value into two pieces:

$$\begin{aligned} \int_{-3/2}^{3/2} \frac{2}{\sqrt{|4u^2-1|}} du &= 2 \int_0^{3/2} \frac{2}{\sqrt{|4u^2-1|}} du = 2 \int_0^{1/2} \frac{2}{\sqrt{1-4u^2}} du + 2 \int_{1/2}^{3/2} \frac{2}{\sqrt{4u^2-1}} du \\ &= \int_0^{\pi/2} \frac{\cos \theta d\theta}{\sqrt{1-\sin^2 \theta}} + \int_{\pi/2}^{\cos^{-1}1/3} \frac{\sec \theta \tan \theta}{\sqrt{\sec^2 \theta - 1}} d\theta \end{aligned}$$

which involve the trig substitutions: $2u = \sin \theta \Rightarrow 2du = \cos \theta d\theta$ and $2u = \sec \theta \Rightarrow 2du = \sec \theta \tan \theta d\theta$

$$\begin{aligned} \int_0^{\pi/2} d\theta + \int_{\pi/2}^{\cos^{-1}1/3} \sec \theta d\theta &= \frac{\pi}{2} + \lim_{b \rightarrow \pi/2^-} \ln |\sec \theta + \tan \theta| \Big|_b^{\cos^{-1}1/3} \\ &= \frac{\pi}{2} + \ln \left| 3 + \frac{\sqrt{8}}{3} \right| - \lim_{b \rightarrow \pi/2^-} \ln |\sec b + \tan b| \\ &= \frac{\pi}{2} + \ln \left| 3 + \frac{\sqrt{8}}{3} \right| + \infty = \infty \end{aligned}$$

